



doi: 10.3978/j.issn.1005-6947.2015.11.027

http://dx.doi.org/10.3978/j.issn.1005-6947.2015.11.027

Chinese Journal of General Surgery, 2015, 24(11):1630-1632.

III

30

733000

			III		
	30	II		III	
		2	6.67%		3
					10.0%
		2	6.67%		
		170 min			15-45
					3 mm
			III		26
	/		/		
	R737.9				
					1.2
					1.3
(1)	(2)				8-10 c
		III	20%		90°
		III		Halsted	
2013	6	-2015	3	30	II
				III	
					1
1					
1.1					
	2013	6	-2015	3	30
II			35-60		II _A
13	II _B	17		43.3	

2015-08-27

2015-10-19

Email: wuweilw@163.com

- -

III

III

2

2.1

30

170 min

2 6.67%

3 10.0%

III

III

III

III

III

2.2

15~45

3 26 III

3~12 6.5 III

21.74% II_A 13

II_B 17

2.3

2015 7 30 III

100% 2 6.67%

1 Rotter

3

[3]

[4]

III

Level-III

[5]

III

II

Auchincloses

Patty [6]

III



doi: 10.3760/cjgs201409020001
http://dx.doi.org/10.3760/cjgs201409020001
Chinese Journal of General Surgery, 2014, 49(9): 100-101



100-101

100

2014年9月2日
第49卷第9期
第100-101页
ISSN 1007-1226
CN 44-1536/R
Y28B

JOURNAL OF GENERAL SURGERY

中华普通外科杂志
Chinese Journal of General Surgery, 2014, 49(9): 100-101
doi: 10.3760/cjgs201409020001
http://dx.doi.org/10.3760/cjgs201409020001

100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120

100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120

100-101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120

Cite this article as:
100-101
doi: 10.3760/cjgs201409020001
http://dx.doi.org/10.3760/cjgs201409020001